



After Your Biopsy

1. Keep the dressing dry for _____ hours. Then you may remove it.
2. After removing the dressing, you may shower. Wash the incision with soap and water, then pat dry.
3. If “Steri-strips” are in place, let them fall off on their own. This usually happens in 7 - 10 days.
4. If Dermabond is in place, refer to the printed instruction sheet about this product.
5. Refrain from strenuous exercise or activity for 1 week.
6. If you have pain, use Tylenol, Motrin, or Advil, if no other pain medication was prescribed.
7. If you have any of the following signs of infection, contact your medical staff fellows:
 - fever
 - chills
 - increased redness and tenderness at the site
 - excessive drainage from the site

Other Instructions:

Contacts (Reach the staff members below through the NIH page operator at **301-496-1211**.)

Your Doctor:



2006

This information is prepared specifically for patients participating in clinical research at the NIH Clinical Center at the National Institutes of Health and is not necessarily applicable to individuals who are patients elsewhere. If you have questions about the information presented here, talk to a member of your healthcare team.

Where applicable, brand names of commercial products are provided only as illustrative examples of acceptable products, and do not imply endorsement by NIH; nor does the fact that a particular brand name product is not identified imply that such product is unsatisfactory.

Questions about the Clinical Center?
OCCC@cc.nih.gov

National Institutes of Health
NIH Clinical Center
Bethesda, Maryland 20892