Procedures/Diagnostic Tests

Después de recibir una transfusión de sangre: Instrucciones para pacientes externos

Usted puede reiniciar sus actividades normales cuatro a seis horas después de la transfusión de sangre. Las reacciones a una transfusión de sangre pueden retrasarse.

- Si tiene fiebre con o sin escalofríos en las próximas 48 horas, es posible que se deba a una reacción retrasada a la transfusión.
- Si observa que su orina está oscura, tiene piel amarilla o se siente más cansado(a) que de costumbre durante las próximas dos semanas, usted podría estar teniendo una reacción retrasada a la transfusión.

Si usted cree que está teniendo una reacción retrasada a la transfusión, llame al personal de atención médica de inmediato.

Nombre: _____

No. Telefónico: _____

This information is prepared specifically for persons taking part in clinical research at the National Institutes of Health Clinical Center and may not apply to patients elsewhere. If you have questions about the information presented here, talk to a member of your health care team.

Products/resources named serve as examples and do not imply endorsement by NIH. The fact that a certain product/resource is not named does not imply that such product/resource is unsatisfactory.

National Institutes of Health Clinical Center
Bethesda, MD 20892
8/1/03

Questions about the Clinical Center?
http://www.cc.nih.gov/comments.shtml