



## 输血后门诊指导

---

您在输血 4 至 6 小时以后便可以恢复正常活动。  
输血后可能会有延迟反应。

- 如果您在未来的 48 小时之内出现发烧、发冷的症状，便可能是产生了对输血的延迟反应。
- 如果您在未来的两周内注意到小便颜色变深、皮肤发黄或异常疲劳，便可能是产生了对输血的延迟反应。

如果您认为自己身上出现了输血延迟反应，请立即打电话和您的护理小组的成员联系。

姓名： \_\_\_\_\_

电话号码： \_\_\_\_\_



8/1/03

This information is prepared specifically for patients participating in clinical research at the Warren Grant Magnuson Clinical Center at the National Institutes of Health and is not necessarily applicable to individuals who are patients elsewhere. If you have questions about the information presented here, talk to a member of your healthcare team.

Questions about the Clinical Center? [OCCC@cc.nih.gov](mailto:OCCC@cc.nih.gov)

*Where applicable, brand names of commercial products are provided only as illustrative examples of acceptable products, and do not imply endorsement by NIH; nor does the fact that a particular brand name product is not identified imply that such product is unsatisfactory.*

National Institutes of Health  
Warren Grant Magnuson Clinical Center  
Bethesda, Maryland 20892