

CCMD Abstracts 1990-2000

1990

Levine SJ, Masur H, Gill VJ, Feuerstein IM, Suffredini AF, Shelhamer JH, Lane HC, Yarchoan R, Ognibene FP. Effect of antipneumocystis prophylaxis on the diagnosis of *Pneumocystis carinii* pneumonia by induced sputum examination in patients infected with the human immunodeficiency virus. Clin Res 1990;38:461A.

De La Cadena RA, Suffredini AF, Kaufman N, Parrillo JE, Colman RW. Activation of the kallikrein-kinin system after endotoxin administration to normal human volunteers. Clin Res 1990;38:346A .

Martich GD, Parker MM, Ognibene FP, Suffredini AF, Cunnion RE. Serial cardiovascular patterns in septic shock: contrasts between candidemia and bacteremia. Clin Res 1990;38:454A.

Danner RL, Suffredini AF, Van Dervort AL, Ceska M, Stuetz PL, Zablotny J, Patterson EA. Detection of interleukin-6 (IL-6) and interleukin-8 (IL-8) during septic shock in humans. Clin Res 1990;38:352A.

Martich GD, Parker MM, Cunnion RE, Suffredini AF. Effects of pentoxifylline or ibuprofen pretreatment on the cardiovascular response of normal humans to endotoxin. Clin Res 1990;38:453A.

Martich GD, Danner RL, Van Dervort AL, Patterson E, Suffredini AF. Priming of neutrophils after intravenous endotoxin administration to normal humans: effects of oral pentoxifylline or ibuprofen. Clin Res 1990;38:306A.

Hoffman AG, Martich GD, Vadas P, Pruzanski W, Shelhamer JH, Suffredini AF. Elevation in serum phospholipase A2 in healthy humans given intravenous endotoxin after pretreatment with anti-inflammatory agents. Clin Res 1990;38:340A.

Hoffman AGD, Ognibene FP, Suffredini AF, Gill V, Lawrence M, Shelhamer JH. Detection of fungal sterols by gas chromatography/mass spectrometry in bronchoalveolar lavage fluid from critically ill patients with aspergillus pneumonia. Am Rev Respir Dis 1990;141:A892.

Hoffman AGD, Lipschik G, Lawrence M, Kovacs J, Ognibene F, Suffredini AF, Masur H, Shelhamer JH. Pulmonary surfactant abnormalities in AIDS patients with pneumocystis pneumonia. FASEB J 1990;4:A1714.

Pruzanski W, Stefanski E, Wilmore DW, Martich GD, Hoffman AGD, Suffredini AF, Vadas P. Sequential activation of TNF phospholipase A2 axis following i.v. endotoxin challenge in human volunteers. FASEB J 1990;4:A2147.

Hoffman WD, Pollack M, Banks SM, Koev LA, Solomon MP, Danner RL, Hill J, Akin GL, Koles NL, Guelde G, Natanson C. Monoclonal anti-endotoxin antibodies as therapy in a canine model of human septic shock . Crit Care Med 1990;18(4):S260.

Natanson C, Hoffman WD, Reilly JM, Danner RL, Parrillo JE. Fluid loading eliminates difference in cardiovascular responses to dopamine and norepinephrine in awake dogs. Crit Care Med 1990;18(4):S173.

Hoffman WD, Danner RL, Koev LA, Banks SM, Elin RJ, Hosseini JM, Walker LD, Dolan DP, Natanson C. Ability of endotoxin and heat stable components of two strains of *E. coli* to produce lethality in canine septic shock. Crit Care Med 1990;18(4):S211.

Van Dervort AL, Danner RL. Interleukin 8 primes human neutrophils for enhanced production of superoxide. Crit Care Med 1990;18(4):S251.

Suffredini AF, Danner RL, Martich GD, Ceska M, Stuetz PL, Patterson E. Detection of neutrophil activating peptide-1/interleukin 8 after the administration of endotoxin to normal humans. Crit Care Med 1990;18(4):S259.

Danner RL, Suffredini AF, Van Dervort AL, Ceska M, Stuetz PL, Zablotny JA. Neutrophil activating peptide-1/interleukin 8 concentrations in human septic shock. Crit Care Med 1990;18(4):S281.

Natanson C, Reilly JM, Hoffman WD, Danner RL, Parrillo JE. Fluid loading eliminates the differences in cardiovascular responses to dopamine and norepinephrine in conscious dogs. Clin Res 1990;38(2):340A.

Hoffman WD, Danner RL, Koev LA, Banks SM, Walker LD, Elin RJ, Hosseini JM, Dolan DP, Natanson C. Ability of endotoxin and heat stable components of two strains of *E. coli* to produce lethality in canine septic shock. Clin Res 1990;38(2):454A.

Hoffman WD, Pollack M, Banks SM, Koev LA, Solomon MA, Danner RL, Koles N, Guelde G, Hill J, Natanson C. Monoclonal anti-endotoxin antibodies as therapy in a canine model of human septic shock. Clin Res 1990;38(2):454A.

Eichacker PQ, Banks SM, Hoffman WD, Richmond SE, Mouginis T, Danner RL, Parrillo JE, MacVittie TJ, Natanson C. Levels of total body oxygen consumption and oxygen extraction do not differ between survivors and nonsurvivors in a canine model of human bacterial sepsis. Clin Res 1990;38(2):260A.

Eichacker PQ, Banks SM, Hoffman WD, Danner RL, Mouginis T, Richmond SE, Farese A, Parrillo JE, MacVittie TJ, Natanson C. Pulmonary injury in a lethal canine model of human bacterial sepsis. Clin Res 1990;38(2):260A.

Van Dervort AL, Danner RL. Superoxide production by human neutrophils is enhanced by interleukin 8. Clin Res 1990;38(2)350A.

Van Dervort AL, Stuetz PL, Danner RL. Lipid X inhibition of endotoxin priming of neutrophils can be changed from competitive to non-competitive by altering the structure. Clin Res 1990;38(2):453A.

Martich GD, Danner RL, Ceska M, Stuetz PL, Patterson E, Suffredini AF. Neutrophil activating peptide-1/interleukin 8 and tumor necrosis factor detection following intravenous endotoxin administration to normal humans. Clin Res 1990;38(2):478A.

Farese AM, Eichacker PQ, Patchen M, Bowles C, Danner RL, Natanson C, MacVittie TJ. Gram negative hyperdynamic sepsis in the canine: systemic and pulmonary neutrophil kinetics and functions. ISEH 1990;18:697.

Farese A, Eichacker P, Patchen M, Eichenholz PW, Natanson C, MacVittie TJ. In vivo infusion of recombinant human F (rhTNF) in normal canines effects the number and function of circulatory neutrophils (DMNS) FASEB J 1990; 4:A524.

Levine SJ, Martinez A, Walsh TJ, Eichacker PQ and Natanson C. Hypoxemia, pulmonary infiltrates, and myocardial depression as a sequelae of liposomal Amphotericin B infusion. American Thoracic Society 1990;141:A600.

Eichacker PQ, Hoffman WD, Mouginis T, Richmond S, MacVittie TJ, and Natanson C. The relationship of oxygen consumption to oxygen delivery differs in anesthetized compared to conscious dogs. Am Rev Resp Dis 1990;141:A584.

Eichacker PQ, Hoffman WD, Kuo G, Richmond S, Banks SM, MacVittie TJ, and Natanson C. Tumor necrosis factor but not Interleukin-1 produces sustained reversible hypoxemia alveolar protein and neutrophil accumulation and hemodynamic depression in a canine model. Am Rev Resp Dis 1990;141:A513.

Eichacker PQ, Banks SM, Alling DW, Hoffman WD, Mouginis T. Richmond S, MacVittie TJ, and Natanson C. The relationship of oxygen consumption to oxygen delivery differs in anesthetized vs. conscious dogs. Crit Care Med 1990;18:5183.

Eichacker, Gruber D, Mouginis T, Hoffman WD, Banks SM, Farese AM, Richmond S, MacVitties TJ, and Natanson C. Antibody against Neutrophil adhesion complex CDW18 does not improve survival or prevent cardiopulmonary abnormalities after tumor necrosis factor infusion in dogs. Crit Care Med 1990;18:5259.

Eichacker PQ, Hoffman WD, Kuo G, Richmond S, Banks SM, MacVittie TJ, and Natanson C. Tumor Necrosis factor but not Interleukin-1 induces sustained but reversible hypoxemia, alveolar protein and neutrophil accumulation and hemodynamic depression in dogs. Crit Care Med 1990;18:S281.

Eichacker PQ, Gruber D, Nouginis T, Hoffman WD, Banks SM, Farese AM, Richmond S, MacVittie TJ, and Natanson C. Antibody against neutrophil adhesion complex CDW11 does not improve survival or prevent cardiopulmonary abnormalities after tumor necrosis factor infusion in dogs. Clin Res 1990;38:260A.

Hoffman, A , Eichacker, P , Lawrence, M, and Shelhammer, JH. Bronchoalveolar lavage lipid profiles in normal dogs. Clin Res 1990 38:572A.

Parker MM, Ognibene FP, Parrillo JE. Reversible depression of myocardial function in septic shock (SS) confirmed by load-independent measure of ventricular performance. Clin Res 1990;38:2:340.

Masur H, Ognibene FP, Suffredini AF, O'Neill D, Gill VJ, Manichewitz J, Stock F, Callaway J, Travis W, Solomon D, Kovacs JA, Davey R, Polis M, Falloon J, Lane HC, Shelhamer JH. Diagnosis of CMV pneumonia in patients with HIV infection. Program and Abstracts of the 30th Interscience Conference on Antimicrobial Agents and Chemotherapy 1990;425.

Levine SJ, Martinez A, Walsh TJ, Eichacker PQ, and Natanson C. Hypoxemia, pulmonary infiltrates, and myocardial depression as a sequelae of liposomal Amphotericin B infusion. Am Rev Resp Dis 1990;141:600.

Eichacker PQ, Hoffman WD, Mouginis T, Richmond S, MacVittie TJ, and Natanson C. The relationship of oxygen consumption to oxygen delivery differs in anesthetized compared to conscious dogs. Am Rev Resp Dis 1990;141:584.

Eichacker PQ, Hoffman WD, Kuo G, Richmond S, Banks SM, MacVittie TJ, and Natanson C. Tumor necrosis factor but not Interleukin-1 produces sustained reversible hypoxemia alveolar protein and neutrophil accumulation and hemodynamic depression in a canine model. Am Rev Resp Dis 1990;141:513.

Eichacker PQ, Banks SM, Alling DW, Hoffman WD, Mouginis T, Richmond S, MacVittie TJ, and Natanson C. The relationship of oxygen consumption to oxygen delivery differs in anesthetized vs. conscious dogs. Crit Care Med 1990;18:5183.

Eichacker, Gruber D, Mouginis T, Hoffman WD, Banks SM, Farese AM, Richmond S, MacVittie TJ, and Natanson C. Antibody against neutrophil adhesion complex CDW18 does not improve survival or prevent cardiopulmonary abnormalities after tumor necrosis factor infusion in dogs. Crit. Care Med 1990;18:5259.

Eichacker PQ, Hoffman WD, Kuo G, Richmond S, Banks SM, MacVittie TJ, and Natanson C. Tumor necrosis factor but not interleukin-1 induces sustained but reversible hypoxemia, alveolar protein and neutrophil accumulation and hemodynamic depression in dogs. Crit Care Med 1990;18:S281.

Eichacker PQ, Gruber D, Mouginis T, Hoffman WD, Banks SM, Farese AM, Richmond S, MacVittie TJ, and Natanson C. Antibody against neutrophil adhesion complex CDW11 does not improve survival or prevent cardiopulmonary abnormalities after tumor necrosis factor infusion in dogs. Clin Res 1990; 38:260.

Farese A, Eichacker P, Patchen M, Eichenholz PW, Natanson C, and MacVittie TJ. In vivo infusion of recombinant human TNF (rhTNF) in normal canines effects the number and function of circulatory neutrophils (DMNS). FASEB 1990;4:524.

Hoffman WD, Koev LA, Solomon MA, Banks SM, Yatsiv I, Dolan D, Clements SL, and Natanson C. Compared to enflurane, isoflurane produces less depression of the stroke volume and autonomic response of canines given phenylephrine. *Anesthesiology* 1990;73:410.

1991

Catania A, Suffredini AF, Lipton JM. Alpha-MSH response to endotoxin in normal human subjects. *FASEB J* 1991;5:A1392 .

Martich GD, Boujoukos AJ, Reda DG, Suffredini AF. Relation of serum neopterin to the hemodynamic and cytokine response following intravenous endotoxin administration to normal humans. *Crit Care Med* 1991 19;S14.

Martich GD, Suffredini AF. Increased oxygen consumption and delivery and widened alveolar-arterial oxygen gradient following intravenous endotoxin administration to normal humans: effects of pretreatment with anti-inflammatory agents. *Am Rev Respir Dis* 1991;143:A204 .

Solomon MA, Correa R, Hoffman WD, Cunnion RE, Cobb JP, Danner RL, Ferrans VJ, and Natanson C. Ultrastructural and histologic changes associated with myocardial depression in a canine model of septic shock. *Clin Res* 1991;39(2):321A.

Eichenholz PW, Eichacker PQ, Hoffman WS, Danner RL, Banks SM, Parrillo JE, Natanson C. Increasing doses of tumor necrosis factor produce different patterns of acute and subacute myocardial dysfunction. *Clin Res* 1991;39(2):321A.

Cobb JP, Natanson C, Solomon MA, Koev LA, Hoffman WD, Banks SM, Danner RL. NG-amino-L-arginine, an inhibitor of nitric oxide synthase, increases lethality in canine endotoxic shock. *Clin Res* 1991;39(2):164A.

Hoffman WD, Pollack M, Banks SM, Koev LA, Solomon MA, Danner RL, Koles N, Guelde G, Yatsiv I, Mouginis T, Elin RJ, Hosseini JM, Porter JC, Natanson C. A controlled trial of IgG monoclonal anti-endotoxin antibodies in a canine model of human septic shock. *Clin Res* 1991;39(2):164A.

Yatsiv I, Koev LA, Hoffman WD, Solomon MA, Danner RL, Dolan D, Banks SM, Natanson C. A controlled trial of continuous arterial-venous hemofiltration fails to improve outcome in a canine model of human septic shock. *Clin Res* 1991; 39(2):264A.

Solomon MA, Alexander HR, Balaban RS, Cobb JP, Hoffman WD, Koev LA, Cunnion RE, Yatsiv I, Banks SM, Kim DK, Danner RL, Roberts WC, Natanson C. Myocardial systolic phosphorylation potential in a canine sepsis model. *Clin Res* 1991;39(2):164A.

Eichacker PQ, Farese A, Hoffman WD, Banks SM, Waisman Y, Danner RL, MacVittie TJ, Natanson C. Alveolar neutrophils are maximal during recovery of pulmonary function in a canine model of human septic shock. *Clin Res* 1991;39(2):453A.

Van Dervort AL, Stuetz P, Danner RL. 3-Aza-lipid X-4-phosphate specifically antagonizes endotoxin-induced priming of human neutrophils independent of systolic calcium. Clin Res 1991;39(2):164A.

Eichacker PQ, Moughinis T, Farese A, Hoffman WD, Banks SM, Suffredini AF, Waisman H, Danner RL, Natanson C. Serum tumor necrosis factor increases early and transiently in a chronic canine model of septic shock. Am Rev Resp Dis 1991;143:A681.

Solomon MA, Kim DK, Balaban RS, Hoffman WD, Koev LA, Alexander HR, Cobb JP, Cunnion RE, Banks SM, Yatsiv I, Danner RL, Roberts WC, Natanson C. Myocardial high energy phosphate stores in a canine model of septic shock. Crit Care Med 1991;S567.

Moughinis T, Natanson C, Hoffman WD, Banks SM, Suffredini AF, Waisman Y, Danner RL, Richmond SE, MacVittie TJ, Eichacker PQ. Serum tumor necrosis factor increases early and transiently in a chronic canine model of septic shock. Crit Care Med 1991;19:S567.

Hoffman WD, Pollack M, Banks SM, Koev LA, Solomon MA, Danner RL, Koles N, Guelde G, Yatsiv I, Moughinis T, Elin RJ, Hosseini JM, Porter JC, Natanson C. Efficacy of monoclonal anti-endotoxin antibodies in canine model of human septic shock. Crit Care Med 1991;19:S167.

Van Dervort AL, Lipschik GY, Stuetz P, Danner RL. A lipid X analog inhibits endotoxin-induced priming without affecting systolic calcium and blocks transcription of the tumor necrosis factor gene. Crit Care Med 1991;19:S57.

Waisman Y, MacVittie T, Farese A, Banks SM, Hoffman WD, Souza L, Natanson C and Eichacker PQ. A controlled trial of granulocyte colony stimulating factor in a canine model of human bacterial sepsis. Clin Res 1991; 39:165A.

Eichacker PQ, Waisman Y, Banks SM, Richmond S, MacVittie T, and Natanson C. Effects of pentothal anesthesia on oxygen delivery and consumption in a canine model. Clin Res 1991;39:265A.

Eichacker PQ, Waisman Y, Banks SM, Richmond S, MacVittie TJ and Natanson C. Effects of pentathol anesthesia on oxygen delivery and consumption in a canine model. Crit Care Med 1991;19:S53.

Waisman Y, Eichacker PQ, Richmond S, Banks SM, Hoffman WD, MacVittie T, and Natanson C. Arterial bases deficit predicts volume loss in canine hemorrhage. Pediatr Res 1991;29(4):35A.

Farese AM, Eichacker PQ, Patchen LM, Natanson C and MacVittie TJ. Gram negative hyperdynamic sepsis in the canine systemic and pulmonary neutrophil kinetics and functions. Second International Congress on the Immune Consequences of Trauma, Shock and Sepsis Mechanisms and Therapeutic Approaches, p76, 1991.

Waisman Y, Eichacker PQ, Richmond S, Banks SM, Hoffman WD, MacVittie T, and Natanson C. Arterial bases deficit predicts volume loss in canine hemorrhage. *Pediatr Res* 1991;29(4):35A.

Levine SJ, Shelhamer JH, Feuerstein IM, Gill VJ, Masur H, Ognibene FP. Utilization of three staining techniques and radiographically directed bronchoalveolar lavage (BAL) results in equivalent diagnostic sensitivity for *Pneumocystis carinii* pneumonia (PCP) in AIDS patients with or without prior aerosolized pentamidine (AP) prophylaxis. *Am Rev Resp Dis* 1991;143:A107.

Ognibene FP, Levine SJ, Shelhamer JH, Feuerstein IM, Gill VJ, Masur H. Diagnostic yield for *Pneumocystis carinii* pneumonia (PCP) in AIDS patients increases with multiple staining techniques and radiographically directed bronchoalveolar lavage (BAL). *Clin Res* 1991;39,2:217.

Orcutt TA, Godwin CR, Pizzo PA, Ognibene FP. Aerosolized pentamidine: prophylaxis against *Pneumocystis carinii* pneumonia (PCP) is well-tolerated in children with AIDS. *Respiratory Care* 1991;36,11:15-16.

Quezado Z, Wilson WH, Parker MM, Cunnion RE, Reda D, Bryant G, Ognibene FP. Ifosfamide induction therapy is associated with severe, reversible cardiac dysfunction. *Clin Res* 1991;39,3:667.

1992

Boujoukos AJ, Martich GD, Supinski E, Suffredini AF. Compartmentalization of acute cytokine responses in humans after intravenous endotoxin administration: characteristics of bronchoalveolar lavage cytokine mRNA and protein. *Am Rev Respir Dis* 1992;145:A441

Boujoukos AJ, Martich GD, Danner RL, Suffredini AF. Detection of interleukin-8 in bronchoalveolar lavage without alveolar neutrophil influx, before and after intravenous endotoxin in normal humans. *Am Rev Respir Dis* 1992;145:A441.

Two Hawk S, Boujoukos AJ, Suffredini AF. High output states in patients with nonHodgkin's lymphoma. *Clin Res* 1992;40:412A.

Martich GD, Van Dervort AL, Danner RL, Suffredini AF. Intravenous endotoxin administration to normal humans primes neutrophils for an enhanced respiratory burst. *Crit Care Med* 1992;20:S100.

Boujoukos AJ, Martich GD, Suffredini AF. Compartmentalization of the acute cytokine response in humans after intravenous endotoxin. 2nd Conference of the International Endotoxin Society. Vienna, Austria Aug 17-20, 1992.

Sylvester I, Suffredini AF, Danner RL, Leonard EJ. Effect of diluents and capture on detection of NAP-1 by sandwich ELISA. 3rd International Meeting of Chemotactic Cytokines. Vienna, Austria. September 1992.

Eichacker PQ, MacVittie TJ, Farese A, Waisman Y, Banks SM, Hoffman WD, Danner RL, Rothlein R, Wilson L, Natanson C. Antibody directed against leukocyte CD18 membrane protein does not improve outcome in a canine model of human sepsis. Am Rev Resp Dis 1992;145:A317.

Eichacker PQ, Hoffman WD, Banks SM, Danner RL, Richmond S, Wilson L, MacVittie TJ, Natanson C. Serial metabolic cart and hemodynamic estimates of total body oxygen consumption in an awake canine model of human septic shock. Am Rev Resp Dis 1992;145:A790.

Cobb JP, Natanson C, Koev CA, Hoffman WD, Griffith OW, Kilbourn RG, Gross S, Levi R, Akin G, Banks SM, Lodato RF and Danner RL. The hemodynamic effects of N -methyl-L-arginine, an inhibitor of nitric oxide synthase, in endotoxemic compared to normal canines. The Society of University Surgeons Residents Program, 1992.

Eichacker PQ, MacVittie TJ, Farese A, Waisman Y, Banks SM, Hoffman WD, Danner RL, Rothlein R, Richmond S, Wilson L, Natanson C. Antibody against leukocyte CD18 membrane protein does not improve outcome in a canine model of human sepsis. Crit Care Med 1992;20:S51.

Cobb JP, Natanson C, Koev CA, Hoffman WD, Griffith OW, Kilbourn RG, Gross S, Levi R, Akin G, Banks SM, Lodato RF and Danner RL. The hemodynamic effects of N -methyl-L-arginine, an inhibitor of nitric oxide synthase, in endotoxemic compared to normal canines. Crit Care Med, 20:S51, 1992.

Quezado ZMN, Natanson C, Banks SM, Alling DW, Koev CA, Danner RL, Elin RJ, Hosseini JM, Pollack M, Bacher JD, Dolan DP, and Hoffman WD. A human IgM monoclonal antibody (MAb) against endotoxin (HA-1A) decreased survival in a canine model of gram-negative bacterial septic shock. Clin Res 1992;20:286A.

Cobb JP, Natanson C, Koev CA, Hoffman WD, Griffith OW, Kilbourn RG, Gross S, Levi R, Akin G, Banks S, Lodato RF and Danner RL. The cardiovascular effects of N -methyl-L-arginine, a nitric oxide synthase inhibitor, in endotoxemic and normal canines. Clin Res 1992;40:267A.

Van Dervort AL, Madara PJ, Cobb JP, Corriveau CC, Tropea MM, Danner RL. The effects of nitric oxide on endotoxin-induced TNF transcription and translation in human neutrophils. Clin Res 1992;40:306A.

Corriveau CC, Cobb JP, Tropea MM, VanDervort AL, Madara PJ, Danner RL. Activation of soluble granylate cyclase by nitric oxide dose not affect human neutrophil chemotaxis. Pediatr Res 1992;31(4):159A.

Quezado ZMN, Natanson C, Banks SM, Alling DA, Koev CA, Danner RL, Elin RJ, Hosseini JM, Bacher JD, Dolan DP, Hoffman WD. Physiologic results from a controlled trial of human IgM monoclonal antibody against lipid A (HA-1A) in a canine model of gram negative septic shock. Interscience Conference on Antimicrobial Agents and Chemotherapy 1992;32:308.

Waisman Y, MacVittie TJ, Natanson C, Farese A, Banks SM, Hoffman WD, Wilson L, Souza L, Eichacker PQ. A controlled trial of pre-treatment with granulocyte colony stimulating factor in a canine model of human bacterial sepsis. Am Rev Resp Dis 1992;145:A417.

Waisman Y, MacVittie TJ, Natanson C, Farese A, Banks SM, Hoffman WD, Wilson L, Souze L, and Eichacker PQ. A controlled trial of treatment with granulocyte colony stimulating factor in a canine model of human bacterial sepsis. Crit Care Med 1992; 20:S99.

Waisman Y, Eichacker PQ, Banks SM, Hoffman WD, MacVittie TJ, and Natanson C. Comparison of prediction models to quantify acute blood loss in canines. Clin Res 1992;20:372A.

Yatsiv I, Quezado ZMN, Hoffman WD, Koev CA, Winkelstein JA, Cork LC, Eichacker PQ, Banks SM and Natanson C. The third component of complement (C3) is necessary for the febrile response after endotoxin challenge in dogs. Clin Res 1992;40:163A.

Sarosy G, Kohn E, Link C, Adamo D, Davis P, Ognibene FP, Goldspiel B, Christian M, Reed E. Taxol dose intensification (DI) in patients with recurrent ovarian cancer. Program/Proceedings of the 28th Annual Meeting of the American Society of Clinical Oncology 1992;11:226.

Wexler L, Thiele C, McClure L, Chanock S, Mertins S, Tsokos M, Avila N, Reynolds J, Ognibene FP, Venzon D, Castleberry R, Casper J, Truitt R, Yanelli J, Schwartzenruber D, Rosenberg S, Horowitz M. Adoptive immunotherapy (AI) of refractory neuroblastoma (NB) with tumor-infiltrating lymphocytes (TIL), interferon-gamma (IFN- γ), and interleukin-2 (IL-2). Program/Proceedings of the 28th Annual Meeting of the American Society of Clinical Oncology. 1992;11:368.

Hoffman WD, Koev LA, Solomon MA, Banks SM, Yatsiv I, Cobb JP, and Natanson C. The consistency of the hemodynamic response of canines to enflurane and isoflurane among subjects and days. IARS 74:5138, 1992.

Waisman Y, Eichacker PQ, Banks SM, Hoffman WD, MacVittie TJ, and Natanson C. Arterial base deficit as an accurate indicator of volume loss in canine hemorrhage. American Academy of Pediatrics, 1992.

Waisman Y, MacVittie TJ, Natanson C, Farese A, Banks SM, Hoffman WD, Wilson L, Souza L, and Eichacker PQ. A controlled trial of pre-treatment with granulocyte colony stimulating factor in a canine model of human bacterial sepsis. Am Rev Resp Dis 1992;145:417.

Hoffman WD, Koev CA, Banks SM, Solomon MA, and Natanson C. Phenylephrine decreases canine cardiac performance by effects on diastolic function. Crit Care Med 1992;20:S72.

Waisman Y, MacVittie TJ, Natanson C, Farese A, Banks SM, Hoffman WD, Wilson L, Souze L, and Eichacker PQ. A controlled trial of treatment with granulocyte colony stimulating factor in a canine model of human bacterial sepsis. Crit Care Med 1992;20:S99.

Hoffman WD, Koev CA, Banks SM, Solomon MA, and Natanson C. Phenylephrine decreases canine cardiac performance by effects on diastolic function. Clin Res 1992;20:372.

1993

Nam MH, Reda D, Boujoukos AJ, Agosti J, Suffredini AF. Recombinant human dimeric tumor necrosis factor receptor: safety and pharmacokinetics in human volunteers. Clin Res 1993;41:249A.

Nam MH, Reda D, Boujoukos AJ, Martich GD, Agosti J, Suffredini AF. Soluble recombinant human interleukin 1 receptor: safety and pharmacokinetics in human volunteers. Clin Res 1993;41:380A.

Nam MH, Miller R, Banks S, Suffredini AF. Effects of recombinant human dimeric tumor necrosis factor receptor on endotoxin-induced whole blood cytokine production. Clin Res 1993;41:139A.

Quezado ZMN, Natanson C, Hoffman WD, Danner RL, Banks S, Dolan D, Hilton A, Eichacker PQ. Cardiopulmonary effects of inhaled nitric oxide in canine lobar and diffuse gram-negative pneumonia at graded levels of oxygen administration. Am Rev Resp Dis 1993;147:720.

Natanson C, Reilly JM, Hoffman WD, Banks SM, Danner RL, Cunnion RE, Parrillo JE. Effect of septic shock on cardiovascular responses to dopamine and norepinephrine in a conscious canine model. Crit Care Med 1993;21:S277.

Quezado ZMN, Natanson C, Banks SM, Koev CA, Danner RL, Elin RJ, Hosseini JM, Bacher JD, Hoffman WD. Pentoxyfylline can increase tumor necrosis factor level and decrease survival in a canine model of gram-negative septic shock. Crit Care Med 1993;21:S281.

Quezado ZMN, Natanson C, Hoffman WD, Banks SM, Danner RL, Koev CA, Eichacker PQ. Cardiopulmonary effects of inhaled nitric oxide in canine lobar and diffuse gram-negative pneumonia. Crit Care Med 1993;21:S210.

Quezado ZMN, Hoffman WD, Banks SM, Alling DW, Koev CA, Danner RL, Elin RJ, Hosseini JM, Bacher JD, Parker TS, Levine DM, Rubin AL, Natanson C. Therapeutic trial of reconstituted human high density lipoprotein in a dog model of gram-negative septic shock: preliminary results. Crit Care Med, 21:S192, 1993.

Cobb JP, Natanson C, Quezado ZMN, Hoffman WD, Koev CA, Banks SM, Elin RF, Hosseini JM, Winkelstein BA, Danner RL. Lack of protection by N^ω-methyl-L-arginine, a nitric oxide synthase inhibitor, in endotoxin challenged canines. Crit Care Med 1993;21:S280.

Cobb JP, Natanson C, Quezado ZMN, Hoffman WD, Koev CA, Banks S, Elin RF, Hosseini JM, Danner RL. Effect of N^ω-methyl-L-arginine (NMA), a nitric oxide synthase inhibitor, on survival in endotoxin (LPS)-challenged canines. Clin Res 1993;41:253A.

Natanson C, Reilly JM, Hoffman WD, Banks SM, Danner RL, Cunnion RE, Parrillo JE. Effect of septic shock on cardiovascular responses to dopamine (DA) and norepinephrine (NE) in a conscious canine model. Clin Res 1993;41:252A.

Quezado ZMN, Natanson C, Banks SM, Loev CA, Danner RL, Elin RJ, Hosseini JM, Bacher JD, Hoffman WD. Pentoxifylline increased tumor necrosis factor level and decreased survival in canine model of gram-negative septic shock. Clin Res 1993;41:252A.

Quezado ZMN, Hoffman WD, Banks SM, Alling DW, Koev CA, Danner RL, Elin RJ, Hosseini JM, Bacher JD, Parker TS, Levine DM, Rubin AL, Natanson C. Therapeutic Trial of reconstituted human high density lipoprotein in a dog model of gram-negative septic shock. Clin Res 1993;41:175A.

Van Dervort AL, Yan L, Madara PJ, Cobb JP, Wesley RA, Tropea MM, Danner RL. Nitric oxide (NO) increases endotoxin (LPS)-induced cytokine production by human neutrophils (PMNs). Clin Res 1993;41:281A.

Yan L, Danner RL. Nitric oxide synthase (NOS) activity in human neutrophils (PMNs). Clin Res 1993;41:247A.

Corriveau CC, Tropea MM, Yan L, Cobb JP, Van Dervort AL, Madara PJ, Danner RL. Effect of nitric oxide on neutrophil chemotaxis and PGE₂ production. Sixth Pediatric Critical Care Colloquium, 39, 1993.

Patterson M, Natanson C, Hoffman WD, Andrawis J, Banks SM, Richmond S, Wilson L, and Eichacker PQ. Pulmonary effects of granulocyte colony stimulating factor treatment during oxygen exposure in non-neutropenic rats. Am Rev Respir Dis 1993;147:96.

Eichacker PQ, Waisman Y, Hoffman WD, Banks SM, Richmond S, Wilson L, and Natanson C. Effects of pentathol anesthesia, paralysis, and mechanical ventilation on oxygen delivery and consumption during graded decreases in blood flow in the canine. Am Rev Respir Dis 1993;147:620.

Quezado ZMN, Hoffman WD, Yatsiv I, Koev CA, Winkelstein JA, Winkelstein BA, Cork LC, Banks SM, Eichacker PQ, and Natanson C. The third component of complement protects against endotoxin-induced organ damage. Crit Care Med 1993;21:S285.

Eichacker PQ, Waisman Y, Hoffman WD, Banks SM, and Natanson C. Effects of pentathol anesthesia, paralysis, and mechanical ventilation on oxygen delivery and consumption with decreases in blood flow in the canine. Crit Care Med 1993;21:S198.

Patterson M, Natanson C, Hoffman WD, Anderson J, Banks SM, and Eichacker PQ. Granulocyte colony stimulating factor treatment during oxygen exposure in non-neutropenic rats. Crit Care Med 1993; 21:S133.

Eichacker PQ, Waisman Y, Natanson C, Farese A, Hoffman WD, Banks SM, Elin R, Hosseini J, and MacVittie TJ (Intro by Masur H). Recombinant granulocyte colony stimulating factor reduces endotoxemia and improves cardiovascular function and survival during bacterial sepsis in non-neutropenic canines. Clin Res 1993;41:240.

Quezado ZMN, Hoffman WD, Yatsiv I, Koev CA, Winkelstein JA, Winklestein BA, Cork LC, Banks SM, Wilson L, Elin RJ, Eichacker PQ, and Natanson C. The third component of complement protects against endotoxin-induced multiple organ failure. Clin Res 1993;41:243.

Cohen OJ, Pantaleo G, Walker R, Ognibene FP, Shelhamer JH, Fauci AS. HIV burden in mononuclear cells from peripheral blood versus bronchoalveolar lavage of the same individuals: a comparative analysis. Clin Res 1993;41,2:248.

Cohen OJ, Pantaleo G, Walker R, Ognibene FP, Shelhamer JH, Fauci AS. Comparative analysis of HIV burden in mononuclear cells (MC) from peripheral blood (PB) versus bronchoalveolar lavage (BAL) of the same individuals. Program/Proceedings of the 9th International Conference on AIDS. 1993;1:62.

Rosenthal CH, Ognibene FP. The child in the adult ICU: a misplaced patient or an opportunity for redesign? Proceedings of the 1993 National Teaching Institute of the American Association of Critical Care Nurses 1993;449-451.

Dichter J, Shelhamer JH, Ognibene FP. Yield of repeat bronchoscopy (FOB) may be dependent on underlying disease. Clin Res 1993;41,3:593.

Eichacker PQ, Waisman Y, Hoffman WD, Banks SM, Richmond S, Wilson L, and Natanson C. Effects of pentathol anesthesia, paralysis, and mechanical ventilation on oxygen delivery and consumption during graded decreases in blood flow in the canine. Am Rev Resp 1993;Dis 147:620.

Quezado ZMN, Hoffman WD, Yatsiv I, Koev CA, Winkelstein JA, Winklestein BA, Cork LC, Banks SM, Eichacker PQ, and Natanson C. The third component of complement protects against endotoxin-induced organ damage. Crit Care Med 1993;21:S285.

Koev CA, Natanson C, Quezado ZMN, Mayeux B, and Hoffman WD. Fentanyl may directly block cardiac β -agonist effects of epinephrine in dogs. Crit Care Med 1993;21:S261.

Eichacker PQ, Hammond W, Farese A, Hoffman WD, Danner RL, Banks SM, MacVittie TJ, Natanson C. Serial serum granulocyte colony stimulating factor levels in canine model of bacterial sepsis. Clin Res 1993;41:240A.

Eichacker PQ, Waisman Y, Hoffman WD, Banks SM, and Natanson C. Effects of pentathol anesthesia, paralysis, and mechanical ventilation on oxygen delivery and consumption with decreases in blood flow in the canine. Crit Care Med 1993;21:S198.

Patterson M, Natanson C, Hoffman WD, Andersen J, Banks SM, and Eichacker PQ. Granulocyte colony stimulating factor treatment during oxygen exposure in non-neutropenic rats. Crit Care Med 1993;21:S133.

Cobb JP, Natanson C, Quezado ZMN, Hoffman WD, Koev CA, Banks SM, Elin RF, Hosseini JM, Winkelstein BA, and Danner RL. Lack of protection by N -methyl-L-arginine, a nitric oxide synthase inhibitor, in endotoxin-challenged canines. Crit Care Med 1993;21:S280.

Eichacker PQ, Waisman Y, Natanson C, Farese A, Hoffman WD, Banks SM, Elin R, Hosseini J, and MacVittie TJ (Intr. by Masur H). Recombinant granulocyte colony stimulating factor reduces endotoxemia and improves cardiovascular function and survival during bacterial sepsis in non-neutropenic canines. Clin Res 1993;41:240.

Koev, CA, Natanson C, Quezado ZMN, Mayeux B, and Hoffman WD. Fentanyl may directly block cardiac-agonist effects of epinephrine in dogs. Clin Res 1993;41:343.

Polis MA, Baird B, Manischewitz J, Jaffe HS, Fisher PE, Walker R, Falloon J, Davey RT, Kovacs J, Lane HC, Masur H. A phase I/II dose escalation trial of (S)-1-[3-hydroxy-2-(phosphonylmethoxy) propyl] cytosine (HPMPc) in HIV-infected persons with CMV. 33rd Interscience Conference on antimicrobial agents and chemotherapy, New Orleans, LA 1993.

Kovacs JA, Baseler R, Dewar R, Vogel S, Davey RT Jr, Falloon J, Polis MA, Walker RE, Stevens R, Salzman NP, Metcalf JA, Masur H, Lane HC.. Immunologic enhancement in HIV-infected patients with intermittent interleukin-2 therapy. 33rd Interscience Conference on antimicrobial agents and chemotherapy, New Orleans, LA 1993.

1994

Suffredini AF, Reda D, Agosti J, Banks S. Effects of recombinant human tumor necrosis factor receptor (TNFR:Fc) on the cardiovascular and cytokine responses of normal volunteers following intravenous endotoxin. Crit Care Med 1994;22:A191.

Suffredini AF, Reda D, Agosti J, Banks S. Antiinflammatory effects of recombinant tumor necrosis factor receptor (TNFR:Fc) in normal humans following intravenous endotoxin. Am J Resp Crit Care 1994;149:A240.

Suffredini AF, Reda D, Agosti J, Banks S. Cardiovascular and cytokine responses of normal humans following intravenous endotoxin and recombinant human tumor necrosis factor receptor (TNFR:Fc). Intensive Care Med 1994;20 (suppl 1):S147.

De La Cadena R, Miller M, Stadnicki A, Tropea M, Agosti J, Colman RW, Suffredini AF. Recombinant tumor necrosis factor receptor (TNFR:Fc) alters endotoxin-induced activation of the contact system and fibrinolytic system in human volunteers. Presented at the American Society of Hematology, Dec 1, 1994, Memphis TN.

Cobb JP, Natanson C, Quezado ZMN, Hoffman WD, Koev CA, Banks S, Elin RF, Hosseini JM, Danner RL. The differential hemodynamic effect of nitric oxide synthase inhibition in normal and endotoxemic canines. *Intensive Care Medicine* 1994;20:S47.

Quezado ZMN, Natanson C, Hoffman WD, Karzai W, Koev C, Danner R, Banks S, Wilson L, Dolan DP, Hilton AT, Madden D, and Eichacker PQ. Cardiopulmonary effects of inhaled nitric oxide in gram-negative bacterial pneumonia in awake canines. *Am J Resp and Crit Care Med* 1994;149:813.

Karzai W, Natanson C, Patterson M, Hoffman W, Danner RL, Correa R, Banks S, Richmond S, Fitz Y, Wilson L, and Eichacker PQ. The pulmonary effects of granulocyte colony stimulating factor (G-CSF) during pneumonia and oxygen exposure in non-neutropenic rats. *Am J Resp and Crit Care Med* 1994;149:241.

Eichacker PQ, Waisman Y, Natanson C, Farese A, Hoffman WD, Wilson L, Banks SM, and MacVittie TJ. Recombinant G-CSF reduces endotoxemia and serum TNF levels and improves survival in a canine model of septic shock. *Crit Care Med* 1994;22:A193.

Eichacker, PQ. A canine model of intraperitoneal bacterial sepsis. *Shock*, 1994;2:S44.

Dichter J, Shelhamer JH, Ognibene FP. The yield of repeat bronchoscopy (BR) in immunocompromised (IC) patients may be dependent on the underlying disease. *Am J Resp and Crit Care Med* 1994;149,4:A935.

Eichacker PQ, Waisman Y, Natanson C, Farese A, Hoffman WD, Wilson L, Banks SM, and MacVittie TJ. Recombinant G-CSF reduces endotoxemia and serum TNF levels and improves survival in a canine model of septic shock. *Crit Care Med* 22:193, 1994.

Preas H, Koev CA, Natanson C, Quezado ZMN, Mayeux B, Banks SM, and Hoffman WD. Fentanyl suppresses epinephrine-induced increases in canine cardiac performance. *Am J Resp Crit Care Med* 1994;149:1059.

Karzai W, Natanson C, Patterson M, Correa R, Hoffman W, Banks S, Richmond S, Fitz Y, Wilson L, Kim HJ, Welty S, and Eichacker PQ. Effects of a murine monoclonal antibody (MAb) against leukocyte CD11b adhesion protein during toxic oxygen exposure in rats. *Am J Resp Crit Care Med* 1994;149:430.

Walker RE, Kelly GG, Kilgariff CE, Woody JN, Faloon J, Polis MA, Davey RT, Kovacs JA, Masur H, Lane HC. A pilot study of the safety and antiviral activity of a chimeric Anti-TNF antibody in HIV infected patients. 36th Interscience Conference on antimicrobial agents and chemotherapy, Orlando, FL October 1994.

Piscitelli SC, Kovacs JA, Vogel S, Metcalf JA, Baseler M, Lane HC. Decline in interleukin-2 concentrations during intravenous infusion in HIV-infected patients. 36th Interscience Conference on antimicrobial agents and chemotherapy, Orlando, FL October 1994.

1995

Preas H, Reda D, Tropea M, Agosti J, Banks S, Suffredini A. The effects of recombinant soluble interleukin-1 receptor on acute inflammatory responses of human endotoxemia. Crit Care Med 1995;23:A162.

Preas H, Godin P, Vandivier RW, Reda D, Jubran A, Tobin M, Suffredini A. The effects of cyclooxygenase inhibition on ventilation in normal and endotoxemic human volunteers. Am J Resp Crit Care Med 1995;151:A319.

Godin P, Fleisher L, Eidsath A, Vandivier RW, Preas H, Buchman T, Suffredini A. Approximate entropy is a more compete descriptor of heart rate variability than standard deviation in human endotoxemia. Am J Resp Crit Care Med 1995;151:A319.

Vandivier RW, Eidsath A, Godin P, Reda D, Leighton S, Preas H, Culpepper S, Banks S, Ryley R, Suffredini A, Danner RL. Exhaled nitric oxide in volunteers administered sodium nitroprusside or endotoxin. Am J Resp Crit Care Med 1995;151:A15.

Quezado ZMN, Natanson C, Hoffman WD, Karzai W, Koev C, Danner RL, Banks SM, Wilson L, and Eichacker PQ. Granulocyte colony stimulating factor (GCSF) treatment at the onset of *E.coli*-peritonitis-induced septic shock in canines. Am J Resp Crit Care Med 1995;151:A447.

Freeman B, Natanson C, Quezado Z, Danner RL, Quezado M, Banks S, Fitz Y, Zeni F, and Eichacker PQ. Prophylactic granulocyte colony stimulating factor reduces fever and prolongs survival but may worsen lung function in a canine model of diffuse *E. coli* pneumonia. Am J Resp and Crit Care Med 1995;151:A319.

Freeman B, Correa R, Natanson C, Karzai W, Patterson M, Hoffman W, Banks S, Fitz Y, Danner RL, and Eichacker PQ. CD11b monoclonal antibody (MAb 1B6) reduces lung injury but shortens survival in antibiotic treated rats during pneumonia. Am J Resp Crit Care Med 1995;151:A763.

Danner RL, Cobb JP, Suffredini, Eichacker PQ, Natanson C. Nitric oxide in sepsis: role in inflammation and shock. Shock 1995;3:A199.

Natanson C, Eichacker PQ, Suffredini A, Danner RL. Selected treatment strategies for septic shock based on proposed mechanisms of pathogenesis. Shock 1995;3:A200.

Suffredini A, Natanson C, Eichacker PQ, Danner RL. Anticytokine therapies in septic shock. Shock 1995;3:A201.

Eichacker PQ, Natanson C, Danner RL, Suffredini A. The neutrophil as a therapeutic target in septic shock. Shock 1995;3:A202.

Vandivier RW, Eidsath A, Godin P, Reda D, Leighton S, Preas H, Culpepper S, Banks S, Ryley R, Suffredini A, Danner RL. Exhaled nitric oxide in volunteers administered endotoxin and ibuprofen. *Endothelium* 1995;3:A446.

Freeman B, Correa R., Natanson C, Karzai W, Quezado ZMN, Patterson M, Hoffman W, Banks S, Fitz Y, Danner RL, Wilson L, Eichacker PQ. Inhibition or augmentation of neutrophil function predictably affects lung injury but not survival during hyperoxia and pneumonia in rats. *Journal of Investigative Medicine* 1995;143:358A.

Quezado ZMN, Natanson C, Hoffman WD, Quezado MJN, Freeman BD, Banks SM, and Eichacker PQ. Granulocyte colony stimulating factor (G-CSF) improves endotoxin clearance but not cardiopulmonary injury in canine endotoxin shock. *Am J Resp and Crit Care Med* 1995;151:A320.

Reisberg D, Cowan K, Tolcher A, Zujewski J, Keegan P, Noone M, Denicoff A, Gossard M, Venzon D, Ognibene F, Goldspiel B, Kohler D, O'Shaughnessy J. 96-hour paclitaxel without and with r-verapamil in patients with metastatic breast cancer previously treated with 3- or 24-hour paclitaxel. Program/Proceedings of the 31st Annual Meeting of the American Society of Clinical Oncology 1995;14:180.

Tavakoli S, Reda D, Stock F, Iwane M, Brown D, Ognibene FP. The utility of sputum induction (SI) for the diagnosis of *P. carinii* (PC) in children. *Journal of Investigative Medicine* 1995;43, 3:499.

1996

Preas HL, Tropea M, Reda D, Vandivier RW, O'Grady NP, Culpepper S, Suffredini AF. Bronchial instillation of endotoxin in normal humans results in local and systemic inflammatory responses. *Am J Resp Crit Care Med* 1996;153:A442.

O'Grady NP, Preas HL, Tropea M, Culpepper S, Reda D, Suffredini AF. Bronchial instillation of endotoxin in normal humans results in localized but not systemic activation of neutrophils. *J Investing Med* 1996;44:256A.

Zeni F, Freeman B, Natanson C, Banks S, and Danner RL. The cardiovascular effects and efficacy of N methyl-L-arginine alone or in combination with epinephrine in a canine peritonitis model of septic shock. In: Program and Abstracts of the 24th Annual Meeting of the French Intensive Care Society, Paris. Abstract 039, 1996.

Zeni F, Quezado M, Fontana J, Natanson C, Correa R, Fitz Y, Danner RL, Richmond S, Banks SM, Eichacker PQ. ICAM-1 directed monoclonal antibody worsens survival during *E. coli* pneumonia in rats. *Am J Resp Crit Care Med* 1996;153:A258.

Eichacker PQ, Quezado M, Zeni F, Correa R, Fontana J, Banks S, Fitz S, Danner RL, Natanson C. G-CSF increases or decreases the relative risk of death dependent on site and severity of infection in rats. *Am J Resp Crit Care Med* 1996;153:A252.

Zeni F, Freeman B, Natanson C, Quezado Z, Fontana J, Danner RL, Quezado, Banks S, Fitz Y, Eichacker PQ. Prophylactic G-CSF significantly improves survival and myocardial function but not lung injury in canine *E. coli* pneumonia and sepsis. Am J Resp Crit Care Med, 153:A838, 1996.

Preas H, Natanson C, Banks SM, and Hoffman WD. Hemodynamic effects of isoproterenol are competitively inhibited by fentanyl administration in canines. Anesth Analg 1996;82:S366.

Zeni F, Freeman B, Natanson C, Banks S, and Danner RL. The cardiovascular effects and efficacy of N- methyl-L-arginine (NMA) alone or in combination with epinephrine (EPI) in a canine peritonitis model of septic shock. In Programs and Abstracts of the 24th Meeting of Intensive Care French Society, Paris, January 17-19, 1996.

Quezado MJN, Quezado ZMN, Natanson C, Zeni F, Fitz Y, Banks S, and Eichacker PQ. G-CSF improves and CD11/18 MAb worsens hemodynamics independently during endotoxemia in canines. Am J Resp Crit Care Med 1996;163:A838.

Zeni F, Freeman B, Natanson C, Quezado Z, Fontana J, Danner RL, Quezado M, Banks S, Fitz Y, and Eichacker PQ. Prophylactic G-CSF significantly improves survival and myocardial function but not lung injury in canine *E. coli* pneumonia and sepsis. Am J Resp Crit Care Med 1996;153:A444.

Fontana J, Zeni F, Correa R, Quezado M, Banks S, Natanson C, Fitz Y, Danner RL, and Eichacker PQ. Inhibition of leukocyte CD11b and endothelial ICAM-1 in *E. coli* pneumonia. Am J Resp Crit Care Med 1996;153:A444.

Quezado MJN, Quezado Z, Natanson C, Zeni F, Fontana J, Correa R, Danner RL, Banks SM, Fitz Y, Richmond S, Dolan D, and Eichacker PQ. Survival with 72 to 168 h of inhaled nitric oxide with varying FiO₂'s during *E. coli* pneumonia in rats. Am J Resp Crit Care Med 1996;153:A391.

Zeni F, Vodovotz Y, Wahl S, Roberts A, Quezado M, Natanson C, Fontana J, Fitz Y, Danner RL, Richmond S, Banks SM, and Eichacker PQ. Controlled trial of intravenous or intrabronchial TGF- 1 alone and in combination during *E. coli* pneumonia in rats. Am J Resp Crit Care Med 1996;153:A697.

Zeni F, Quezado M, Fontana J, Natanson C, Correa R, Fitz Y, Danner RL, Richmond S, Banks SM, and Eichacker PQ. ICAM-1 directed monoclonal antibody worsens survival during *E. coli* pneumonia in rats. Am J Resp Crit Care Med 1996;153:A258.

Eichacker PQ, Quezado M, Zeni F, Correa R, Fontana J, Banks S, Fitz Y, Danner RL, and Natanson C. G-CSF increases or decreases the relative risk of death dependent on site and severity of infection in rats. Am J Resp Crit Care Med 1996;153:A252.

Sevransky J, Shaked G, Novogrodsky A, Levitzki A, Gazit A, Quezado Z, Freeman B, Eichacker PQ, Danner RL, Banks SM, and Natanson C. Tyrphostin AG 556 improves survival and reduces multiorgan failure in canine *E. coli* peritonitis. In programs and abstracts of the Molecular Basis of Sepsis Meeting, Woodshole, Boston, October 8-11, 1996.

Freeman B, Zeni F, Natanson C, Banks S, and Danner RL. N- ω -methyl-L-arginine (L-NMMA) alone and in combination with epinephrine (EPI) in a canine peritonitis model of septic shock. In Programs and Abstracts of the Molecular Basis of Sepsis Meeting, Woodshole, Boston, October 8-11, 1996.

Highbarger HC, Jiang MK, Kovacs JA, Vogel S, Lebovics R, Dailey P, Lane HC and Dewar RL. Evaluation of IL-2-induced changes in viral burden in lymphoid tissue. 3rd Conference of on Retroviruses and Opportunistic Infections, Washington, DC February 1996.

Walker RE, Bechtel CM, Natarajan V, Baseler M, Metcalf JA, Stevens R, Blaese RM, Davey RT, Falloon J, Polis MA, Kovacs JA, Masur H, Broad DF, Finer MH, Roberts MR, Otten G, Lane HC. 3rd Conference of on Retroviruses and Opportunistic Infections, Washington, DC February 1996.

Walworth CM, Kovacs JA, Vogel SE, Premkumar A, Feuerstein IM. Factors associated with symptomatic interleukin-2 induced cholestopathy in HIV-infected individuals. 36th Interscience Conference on antimicrobial agents and chemotherapy, New Orleans, LA September 1996.

Benfield T, Lundgren B, Lundgren JD, Shelhamer J. MSG of *P. carinii* induces gene expression and secretion of IL-6 and 8 in alveolar epithelium. Am J Resp Crit Care Med 1996;155(4) .

Logun C, Tavakoli S, Levine S, Benfield T, Shelhamer JH. Induction of interleukin-6 production from the airway epithelium in response to lipid mediators. Am J Resp Crit Care Med 155(4) .

Tavakoli S, Logun C, Levine S, Benfield T, Shelhamer JH. Prostaglandin E2 induces the release of interleukin-6 from cultured airway epithelial cells. Am J Resp Crit Care Med 1996;155(4).

1997

Suffredini AF, Preas HL, O'Grady NP. Endotoxin and the human systemic inflammatory response. Shock (Suppl) 1997;7:2.

O'Grady NP, Preas HL, Vandivier RW, Tropea M, Reda D, Suffredini AF. Macrophage inflammatory peptide (MIP-1 alpha) and (MIP-1 beta) elevations in septic shock and effects of cyclooxygenase or TNF inhibition during endotoxemia. American Thoracic Society Meeting, San Francisco, CA May 1997.

Pugin J, Widmer MC, Kossodo S, Liang C-M, Preas HL, Suffredini AF. Human neutrophils secrete gelatinase B in vitro and in vivo in response to endotoxin and proinflammatory mediators. American Thoracic Society Meeting, San Francisco, CA May 1997.

Esposito C, Zeni F, Danner RL, Suffredini AF, Roettinger A, Quezado M, Natanson C, Banks S, Fitz Y, Eichacker PQ. TNFR:Fc and G-CSF have either opposite or similar effects on survival depending on the site and lethality of infection in rats. American Thoracic Society Meeting, San Francisco, CA May 1997.

Jubran A, Tobin MJ, Preas HL, Eidsath A, Vandivier RW, Godin P, Reda D, Suffredini AF. Effect of endotoxemia on variational activity of breathing in humans. American Thoracic Society Meeting, San Francisco, CA May 1997.

Danner RL, Wang S, Vandivier RW, Freeman BD. The dual nature of nitric oxide in SIRS. Shock, 7(Suppl):2, 1997.

Sevransky JE, Shaked G, Novogrodsky A, Levitzki A, Gazit A, Hoffman A, Quezado Z, Freeman BD, Eichacker PQ, Danner RL, Banks SM, and Natanson C. Tyrphostin AG 556 improves survival and reduces multiorgan failure in canine *E. coli* peritonitis. Crit Care Med 1997;25(1):A27.

Sevransky JE, Shaked G, Novogrodsky A, Levitzki A, Gazit A, Hoffman A, Quezado Z, Freeman BD, Eichacker PQ, Danner RL, Banks SM, and Natanson C. Tyrphostin AG 556 improves survival and reduces TNF levels without affecting endotoxemia or bacteremia in canine *E.coli* peritonitis. Am J Resp and Crit Care Med 1997; 155(4):A264.

Esposito C, Zeni F, Danner RL, Suffredini A, Roettinger A, Quezado M, Natanson C, Banks S, Fitz Y, and Eichacker PQ. TNFR:Fc and G-CSF have either opposite or similar effects on survival depending on the site and lethality of infection in rats. Am J Resp and Crit Care Med 1997;155(4):A705.

Eichacker P. Neutrophil stimulation with G-CSF in small and large animal models of sepsis. Shock 1997; 7 (Suppl), 81.

Eichacker P. The relevance of animal models in sepsis. Proceedings and Abstracts of the Second International Symposium on Infections in the Critically Ill Patient. Barcelona, Spain. pp 43, 1997.

Zeni F, Freeman B, Natanson C, Danner RL, Quezado M, Banks S, Fitz Y, J. Bacher, and Eichacker PQ. Prophylactic rG-CSF in a canine model of *E. coli* pneumonia: a controlled trial. Rean Urg 1997; 5:747.

Zeni F, Fontana J, Natanson C, Correa R, Danner RL, Fitz Y, Banks S, Richmond S, Eichacker PQ. Inhibition of leukocyte CD11b and endothelial ICAM-1 in *E. coli* pneumonia. Rean Urg 1997; 5:747.

Zeni F, Quezado M, Natanson C, Correa R, Fontana J, Banks S, Fitz Y, Danner RL, and Eichacker PQ. G-CSF increases or decreases the relative risk of death dependent on site and severity of infection in rats. *Rean Urg* 1997;5:822.

Zeni F, Freeman B, Banks S, Suffredini A, Eichacker PQ, Danner RL, Dhainaut JF, Natanson C. Clinical trials of nonsteroid anti-inflammatory agents in severe sepsis;a meta-analysis and comparison to high dose glucocorticoid and antiendotoxin clinical trials. *Rean Urg* 1997;5:753.

Medin DL, Brown DT, Ognibene FP, Cunnion RE. Comparison of cardiac output measurements by bolus thermodilution technique and continuous automated thermal technique in critically ill patients. *Crit Care Med* 1997;25:A81 (Suppl).

Slonim AD, Ognibene FP. The use of amnesia agents in pediatric bronchoscopy. *Crit Care Med* 1997;25:A85 (Suppl).

Wichman A, Ognibene FP, Sandler A. Research involving critically ill subjects in emergency circumstances: new regulations, new challenges. 1997 Annual Meeting of the American Association of Bioethics, the Society for Bioethics Consultation, and the Society for Health and Human Values.

Sevransky JE, Shaked G, Novogrodsky A, Levitzki A, Gazit A, Quezado Z, Freeman BD, Eichacker PQ, Danner RL, Banks SM, and Natanson C. Tyrphostin AG 556 improves survival and reduces multiorgan failure in canine *E. coli* peritonitis. *Crit Care Med* 1997;25:A27.

Sevransky JE, Shaked G, Novogrodsky A, Levitzki A, Gazit A, Hoffman A, Quezado Z, Freeman BD, Eichacker PQ, Danner RL, Banks SM, and Natanson C. Tyrphostin AG 556 improves survival and reduces TNF levels without affecting endotoxemia or bacteremia in canine *E. coli* peritonitis. *Am J Resp Crit Care Med* 1997;155:A705.

Walker RE, Hahn B, Davey GG, Falloon J, Kovacs JA, Polis MA, Masur H, Metcalf JA, Baseler M, Fyfe G, Thomas S, McCloskey RV, Lane HC. Effects of TNF-alpha antagonists Thalidomide and monoclonal Anti-TNF (cA2) on reducing IL-2-associated toxicities: a randomized, controlled trial. 5th Conference of on Retroviruses and Opportunistic Infections, Washington, DC February 1997.

Masur, H. Therapy and prophylaxis: why do patients fail and what's ahead? 5th Conference of on Retroviruses and Opportunistic Infections, Washington, DC February 1997.

Yao XL, Wu T, Cowan M, Logun C, Shelhamer JH. The effect of dexamethasone on cytosolic phospholipase A2 gene expression in hela cells. *Am J Resp Crit Care Med* 1997;155(4) .

Cowan MJ, Yao XL, Wu T, Shelhamer JH. Characterization of the promoter of the human cytosolic phospholipase A2 gene. *Am J Resp Crit Care Med* 1997;155(4).

Christman BW, Preas HL, Vandivier RW, Price P, Suffredini AF. Ibuprofen increases lipid peroxidation during human endotoxemia. American Thoracic Society Meeting, Chicago, April 1998.

Parent C, Kalil A, Correa R, Natanson C, Obeid J, Danner RL, Fitz Y, Richmond S, Banks SM, Gerstenberger EP, Lee A, and Eichacker PQ. L-selectin directed MAbs increases or decreases survival dependent on site but not severity of infection in rats. Crit Care Med 1998;26:A30.

Wang W, Wang S, Madara P, Yan L, and Danner RL. Human endothelial nitric oxide synthase (eNOS) regulates TNF production in transfected U937 cells via a reactive oxygen species (ROS) signaling pathway. Journal of Investigative Medicine 1998;46:234A.

Wang S, Wang W, Wesley R, and Danner RL. Nitric oxide activates the tumor necrosis factor- α promoter through an Sp1-binding site. FASEB J 1998;12:A301.

Masur H, Kovacs JA. Cloning and sequencing of the dihydrofolate reductase (DHFR) gene of human *Pneumocystis carinii*. Infectious Disease Society of America, Denver, CO November 1998.

Masur H, Brosgart C, Cooper R, Eron FJ, Feinberg J, Falanigan T, Hammer S, Murphy R, Torres R, Thomas D, Rogers M, Ruiz N, Manion D. Salvage therapy with abacavir, Amprenavir, and Efavirenz in subjects with plasma HIV-1 RNA >500 copies/mL despite protease inhibitor (PI) therapy. Infectious Disease Society of America, Denver, CO November 1998.

Piscitelli S, Vogel S, Sadler B, Fiske W, Metcalf J, Masur H, Falloon J. Effect of efavirenz (DMP 266) on the pharmacokinetics of 141W94 in HIV-infected patients. 5th Conference of on Retroviruses and Opportunistic Infections, Chicago, IL February 1998.

Polis MA, Schrager L, Yoder C, Mican J, Kelly G, Metcalf J, Kovacs J, Walker R, Masur H, Falloon J, Schwartzentruber D, Davey R, Haase A, Fox C, Feinberg M. Virologic, immunologic and histologic responses to a 4-drug combination therapy in antiretroviral naive, HIV infected persons. 5th Conference of on Retroviruses and Opportunistic Infections, Chicago, IL February 1998.

Ma L, Masur H, Kovacs JA. Cloning and sequencing of the dihydrofolate reductase (DHFR) gene of human *pneumocystis carinii*. Abstracts of the IDSA 36th Annual Meeting, Denver, 1998.

Huang SN, Fischer SH, O'Shaughnessy E, Gill VJ, Masur H, Kovacs JA. Diagnosis of *Pneumocystis carinii* Pneumonia (PCP) by PCR using major surface glycoprotein (MSG) gene primers in bronchoalveolar lavage and oropharyngeal samples. Abstracts of the IDSA 36th Annual Meeting, Denver, 1998.

Sumida S, Miller K, Vogel S, Kovacs, J, Falloon J, Davey R, Walker R, Masur H, Lane HC, Polis MA. Hypothyroidism is associated with IL-2 therapy in a randomized controlled trial (RCT) of IL-2 for the treatment of HIV-infection. Abstracts of the IDSA 36th Annual Meeting, Denver, 1998.

Eron J, Falloon J, Masur H, Ait-Khaled M, Thomas D, Manion D, Rogers M. Activity of combination abacavir/amprenavir/efavirenz therapy in HIV-1 infected subjects failing their current protease inhibitor containing regimen. Aids 1998;12:S13-S13.

Huang S, Angus W, Turner R, Sorial V, Kovacs JA. Identification and characterization of a novel major surface glycoprotein like gene family in rat *Pneumocystis carinii*. FASEB Journal 1998;12(8):A1349.

Angus CW, Klivington D, Wyman J, Kovacs JA. Nucleic acid vaccination against *Toxoplasma gondii* in mice. FASEB Journal 1998;12(8):A1399.

Russian DA, Andrawis-Sorial V, Vogel P, Turner RE, Klivington DL, Angus CW, Kovacs JA. A surface expressed serine protease is encoded by a multicopy gene family in *Pneumocystis carinii*. FASEB Journal 1998;12(8):A1432.

1999

O'Grady NP, Tropea M, Preas HL, Reda D, Suffredini AF. Factors delaying apoptosis of elicited neutrophils in humans challenged with endobronchial endotoxin. Crit Care Med 1999; 27: A41 Suppl.

Wang W, Wang S, Madara P, Del Pilar Cintron A, Danner RL. Endothelial nitric oxide synthase upregulates TNF production in transfected U937 cells through superoxide-dependent activation of P44/42 mitogen activated protein kinase. J Invest Med 1999; 47: 213A.

Orenstein J, Feinberg M, Yoder C, Schrager L, Mican J, Metcalf J, Fox C, Masur H. Persistence of follicular hyperplasia in previously antiretroviral naïve HIV-Infected patients After 6 months on aggressive therapy. The 6th Conference on Retroviruses and Opportunistic Infections, Chicago, 1999.

Imamichi H, Zhang YM, Kovacs J, Dewar R, Metcalf J, Vogel S, Salzman N, Lane HC. HIV-1 induced in the setting of IL-2 therapy is transient and not derived from PGMC proviral reservoirs. The 6th Conference on Retroviruses and Opportunistic Infections, Chicago, 1999.

Chun T-W, Engel D, Mizell S, Metcalf J, Hallahan C, Kovacs J, Davey R, Dybul M, Mican J, Lane C, Fauci A. Effect of interleukin-2 in diminution of a pool of latently infected, resting CD4+ T cells in HIV-1 patients receiving highly active antiretroviral therapy. The 6th Conference on Retroviruses and Opportunistic Infections, Chicago, 1999.

Sevransky JE, Vandivier RW, Quezado ZMN, Eichacker PQ, Danner RL, Banks SM, Backer J, Thomas ML, Natanson C. The effect of L-NMMA on TNF induced cardiac dysfunction. Am J Respir Crit Care Med 1999;159 (3):A245.

Parent C, Natanson C, Cui X, Banks S, Eichacker PQ. Control mortality and exclusion criteria: factors potentially altering the effects of anti-inflammatory therapies in clinical sepsis trials. Am J Respir Crit Care Med 1999;159 (3):A263.

Gladwin MT, Yao XL, Cowan MJ, Logan C, Huang XL, Shelhamer JH. Reduction in p11 protein levels by retinoic acid (RA) increases CPLA₂ activity and enhances susceptibility to oxidative cell death. Am J Respir Crit Care Med 1999;159 (3):A273.

Yao XL, Cowan MJ, Gladwin MT, Lawrence M, Shelhamer JH. Dexamethasone inhibits cytosolic phospholipase A₂ activity in human epithelial cells by increasing p11 production. Am J Respir Crit Care Med 1999;159 (3):A333.

Huang XL, Yao XL, Gladwin MT, Cowan MJ, Logun C, Shelhamer JH. Arachidonic acid induces p11 protein expression in human bronchial epithelial cells. Am J Respir Crit Care Med 1999;159 (3):A402.

Cowan MJ, Wu T, Yao XL, Logun C, Shelhamer JH. Initial description of the promotor for the human p11 gene. Am J Respir Crit Care Med 1999;159 (3):A445.

Gladwin MT, Yao XL, Cowan MJ, Logun C, Huang XL, Shelhamer JH. Retinoic acid (RA) reduces p11 protein levels in a human bronchial cell line (BEAS-2B) by a post-translational mechanism. Am J Respir Crit Care Med 1999;159 (3):A447.

Yao XL, Huang XL, Gladwin MT, Cowan MJ, Logun C, Shelhamer JH. Interferon gamma and tumor necrosis factor alpha induce p11 producti8on in human airway epithelial cells. Am J Respir Crit Care Med 1999;159 (3):A448.

DePietro M, Correa R, Zeni F, Vodovotz Y, Natanson C, Wahl S, Roberts A, Fitz Y, Danner R, Banks S, Eichacker PQ. TGF-beta alters survival rates and lung injury dependent on the mortality of *E. coli* challenge in a rat pneumonia model. Am J Respir Crit Care Med 1999;159 (3):A597.

O'Grady NP, Preas HL, Pugin J, Martin TR, Suffredini AF. Inflammatory modulating proteins are found 48 hours after local endotoxin administration in humans. Am J Respir Crit Care Med 1999;159 (3):A601.

Parent C, Esposito C, Kalil A, Natanson C, Sevransky J, Danner R, Banks S, Roettinger A, Obeid J, Fitz Y, Eichacker PQ. Anti-inflammatory agents increase or decrease survival dependent on the severity of infectious challenge in sepsis models. Am J Respir Crit Care Med 1999;159 (3):A614.

Jubran A, Preas HL, Suffredini AF, Tobin MJ, Hines VA. Is variational activity of breathing during endotoxemia in humans modulated by the cyclooxygenase pathway? Am J Respir Crit Care Med 1999;159 (3):A788.

Sevransky JE, Vandivier RW, Quezado ZMN, Eichacker PQ, Danner RL, Banks SM, Bacher J, Thomas ML, Natanson C. The effects of L-NMMA on TNF induced cardiac dysfunction. Am J Respir Crit Care Med 1999; 159: A245.

O'Grady NP, Tropea M, Filie A, Reda D, Suffredini AF. Alterations in lymphocytes during the acute pulmonary inflammatory response to endotoxin in humans. Am J Respir Crit Care Med 1999;159:A601.

Gladwin MT, Ognibene FP, Shelhamer JH, conway DA, Hrinczenko BW, Pannell LK, Nichols JS, Link ME, Pease-Fye ME, Suffredini AF, Noguchi CT, Rodgers GP, Schechter AN. Effects of nitric oxide inhalation on erythrocytes of normal and sickle cell individuals. 1999 Sickle Cell Disease Program. Presented 1999.

Karzai W, Eichacker PQ, Mehlhorn B, Bloos F, Reinhart K. The role of neutrophils in ventilation induced lung injury. Anesthesiology 1999;91(3A):U531.

Kovacs JA, Jackson HL, Schweizer D, Taylor W, Shearer J. Influence that ligand constraints have on the reactivity of synthetic nitrile hydratase models. Abstr Pap Am Chem 1999;218:U792.

Kovacs JA, Taylor WD. Reactivity of a five-coordinate model for nitrile hydratase. Abstracts of Papers of the American Chemical Society 1999;218:U858.

Wang WH, Wang SB, Madara P, Cintron AD, Danner RL. Endothelial nitric oxide synthase upregulates TNF alpha production in transfected U937 cells through superoxide- dependent activation of P44/42 mitogen activated protein kinase. FASEB Journal 1999;13(4):A132.

Lempicki RA, Kovacs JA, Baseler MW, Adelsberger JW, Metcalf JA, Dimitrov DS, Stevens RA, Lambert LA, Alvord WG, Lane HC. Impact of therapy on the kinetics of CD4(+) and CD8(+) T cell turnover in HIV-infected patients receiving HAART or IL-2. FASEB Journal 1999;13(4):A294.

Schweitzer D, Taylor WD, Kovacs JA. Synthetic models of the active site of nitrile hydratase. J Inorg Biochem 1999;74(1-4):291-291.

O'Grady NP, Tropea M, Preas HL, Reda D, Suffredini AF. Factors delaying apoptosis of elicited neutrophils in humans challenged with endobronchial endotoxin. Critical Care Medicine 1999;27(1):A41.

Gladwin MT, Ognibene FP, Shelhamer JH, Pannell LK, Nichols JS, Pease-Fye ME, Schechter AN, The role of hemoglobin in the transport and metabolism of nitric oxide in normal individuals. Blood 1999;94 (Suppl):201A.

Gladwin MT, Ognibene FP, Shelhamer JH, Hrinczenko BW, Pannell LK, Nichols JS, Pease-Fye ME, Noguchi CT, Rodgers GP, Schechter AN. Nitric oxide (NO) inhalation augments nitrosyl (heme) hemoglobin and SNO-hemoglobin formation in SS patients without changing oxygen affinity. Blood 1999; 94 (Suppl):201A.

Preas HL, Becker KL, Snider RH, Nylen ES, White JC, Agosti JM, Suffredini AF. Serum levels of calcitonin precursors are altered by antiinflammatory therapies during experimental endotoxemia. 39th Interscience Conference on Antimicrobial Agents and Chemotherapy, San Francisco, CA. Sept 6-29, 1999, #272, p369.

Fiuza C, O'Grady NP, Tropea M, Reda D, Suffredini AF. Expression of cell surface markers on lavage and blood cells following local pulmonary endotoxin challenge in humans. American College of Surgeons, San Francisco, CA. Oct. 11, 1999.

2000

Paul R, Shelhamer JH, Ognibene FP, Pease-Fye ME, Schechter AN, Gladwin MT. S-nitrosohemoglobin (SNO-Hb) formation by nitric oxide (NO) donors in intact sickle cell erythrocytes. Am J Resp Crit Care Med 2000;161:A848.

Gladwin MT, Shelhamer JH, Pannell LK, Nichols JS, Pease-Fye ME, Schechter AN and Ognibene FP. The role of hemoglobin in the transport and metabolism of nitric oxide in the human circulation. Am J Resp Crit Care Med 2000;161:A848.

Gustavsen KA, Ognibene FP, Reda D, Gill VJ, Lipschik GY. PCR for *Pneumocystis carinii* in patients with Wegener's granulomatosis. Am J Resp Crit Care Med 2000;161:A726.

Gladwin M, Ognibene F, Pease-Fye M, Cannon R, Shelhamer J, Schechter A. Relative role of heme nitrosylation and beta-cysteine 93 nitrosation in the transport and metabolism of nitric oxide by hemoglobin in the human circulation. Nitric Oxide. 2000;4:183.

Cui X, Parent C, Natanson C, Banks S, Gerstenberger E, Fitz Y, Danner RL, Eichacker PQ. The outcome of conventional and new sepsis therapies is determined by the site and dose of bacterial challenge. Am J Resp Crit Care Med 2000;161:A878.

Submitted

Gladwin MT, Shelhamer JH, Schechter AN, Pease-Fye ME, Waclawiw MA, Panza JA, Ognibene FP, Cannon RO. Role of circulating nitrite and S-nitrosohemoglobin in regulating vascular blood flow in man. Submitted.

Cui X, Parent C, Banks S, Gerstenberger E, Fitz Y, Danner RL, Natanson C, Eichacker PQ. TNF soluble receptor (TNFCR:Rc) combined with fluid therapy reduces survival compared to its use alone in septic rats. Submitted.

Alexander E, Susla GM, Burstein AH, Brown D, Ognibene FP. Retrospective evaluation of commonly used equations to predict energy expenditure in mechanically ventilated critically ill patients. Submitted.