Procedures/Diagnostic Tests

Après votre transfusion sanguine: Instructions pour patients en consultation externe

Vous pouvez reprendre vos activités normales au bout de 4 à 6 heures après votre transfusion sanguine. Les réactions suite à une transfusion sanguine peuvent être retardées.

- Si vous développez de la fièvre, avec ou sans frissons, au cours des prochaines 48 heures, il se peut que vous subissiez une réaction tardive à la transfusion.
- Si vous remarquez une urine sombre, une peau jaunâtre ou si vous vous sentez anormalement fatigué(e) au cours des 2 semaines qui suivent, il se peut que vous subissiez une réaction tardive à la transfusion.

Si vous pensez subir une réaction tardive à la transfusion, contactez immédiatement un membre de votre équipe de soins de santé.

Nom: _____

Numéro de téléphone:_____

This information is prepared specifically for persons taking part in clinical research at the National Institutes of Health Clinical Center and may not apply to patients elsewhere. If you have questions about the information presented here, talk to a member of your health care team.

Products/resources named serve as examples and do not imply endorsement by NIH. The fact that a certain product/resource is not named does not imply that such product/resource is unsatisfactory.

National Institutes of Health Clinical Center
Bethesda, MD 20892
8/1/0/03

Questions about the Clinical Center?
http://www.cc.nih.gov/comments.shtml